

Tihomir Vuković, Božina Ivanović and Borivoj Knežević²

**MORPHOLOGICAL ANALYSIS OF SOME HYBRID FISHES
FORMS FROM LAKE SKADAR¹**

MORFOLOŠKA ANALIZA NEKIH HIBRIDNIH OBLIKA RIBA IZ
SKADARSKOG JEZERA

Abstract

During the period from 1971. through 1973. the natural hybrids between the different species from the family of Cyprinidae have been collected in Lake Skadar. The analysis of some meristic and morphometric characters, which is explained in this study, indicates that the collected specimens are hybridized between *Alburnus alburnus alborella* and *Leuciscus cephalus albus*.

Izvod

U periodu od 1971. do 1973. u Skadarskom jezeru su prikupljeni prirodni hibridi među različitim vrstama iz familije Cyprinidae. Analiza nekih merističkih i morfometrijskih karaktera koja je prikazana u ovom radu pokazuje da su prikupljeni primjerci hibridi između *Alburnus alburnus alborella* i *Leuciscus cephalus albus*.

INTRODUCTION

There is no information in the literature about the existence of hybrids between the different fish species from Lake Skadar. Ivanović, B. (1967) submits the results of the experimental

¹ This paper was presented on the I Congressus Europaeus inchnthologorum, Sarajevo (21—29. IX 1973).

² Tihomir Vuković, Biological institute, Sarajevo; Božina Ivanović, Borivoj Knežević, Biological station, Titograd.

hybridization between *Pachychilon pictum* and *Rutilus rubilio* but such natural hybrids have been not yet found in Lake Skadar. Our attention was mainly directed to the species from the family of Cyprinidae, which is represented in Lake Skadar with the large number of species (19). In spite of such a great number of species, it was not described anyone natural hybrid between the species from the family of Cyprinidae. It may be expected that the future study of natural interspecies hybridization of fish hybrid forms which so far have not been recorded in the scientific literature.

MATERIAL AND METHODS

The material has been collected during 1971, 1972. and 1973 from the different localities from lake Skadar near Vranjina. The fish was caught with gill nets with different mesh size. There were caught 11 hybrids in total. The hybrid forms are preserved in 4% formaline for the future investigations. The meristic and morphomeric characters were determined on the base of common schemes (Vuković, T. and Ivanović, B., 1971).

RESULTS AND DISCUSSION

In this preliminary notification was first description of the hybrids *Alburnus alburnus alborella* X *Leuciscus cephalus albus* from Lake Skadar (table 1 and 2).

Tabela 1.

Neki morfološki karakteri hibrida *Alburnus alburnus alborella* (De Filippi, 1844) X *Leuciscus cephalus albus* (Bonaparte, 1838) iz Skadarskog jezera

Table 1.

Some morphological characters of hybrids *Alburnus alburnus alborella* (De Filippi, 1844) X *Leuciscus cephalus albus* (Bonaparte, 1838) from lake Skadar

Mark	Charakters	Amplituda (min.—max.)	M±m(M)	δ
	Body weight, in gr.	271,0—795,0	47,660 ± 5,497	16,492
ab	Total body length, in mm.	127,3—190,0	162,120 ± 6,081	18,245
ad	Body length without c, in	107,5—163,0	137,110 ± 5,283	15,850
In % of body length without c				
ao/ad	Length of head	28,5— 40,7	23,93 ± 0,437	1,313
aq/ad	Anterdorsal distance	59,1— 90,0	54,29 ± 0,592	1,778
rd/ad	Postdorsal distance	39,3— 58,3	36,86 ± 0,412	1,236

Mark	Charakters	Amplituda (min.—max.)	M ± m(M)	δ
lm/ad	Depth of head at occiput	18,2— 27,1	15,79 ± 0,307	0,922
gh/ad	Depth of body	23,9— 38,6	22,08 ± 0,530	1,591
ik/ad	Depth of body on caudal penducle	11,1— 15,7	9,57 ± 0,176	0,529
fd/ad	Lenght of the caudal penducle	21,0— 32,0	20,96 ± 0,375	1,123
vz/ad	Distance between the pectoral and ventral fins	30,9— 41,0	25,70 ± 0,351	1,053
zy/ad	Distance between the ventral and anal fins	24,9— 32,2	21,25 ± 0,318	0,954
gs/ad	Lenght of the base of the dorsal fin	10,6— 18,1	10,74 ± 0,229	0,688
tu/ad	Depth of the anterior part of the dorsal fin	15,7— 25,0	15,30 ± 0,344	1,034
yy/ad	Lenght of the bases of the anal fin	10,1— 24,9	12,15 ± 0,619	1,858
ej/ad	Depth of the anal fin	14,0— 19,7	11,65 ± 0,322	0,966
vx/ad	Lenght of the pectoral fin	17,8— 27,9	17,75 ± 0,235	0,706
zz/ad	Lenght of the ventral fin	14,7— 23,0	14,34 ± 0,331	0,995
In % lenght of head				
np/ao	Diametar of eye	6,0— 7,8	21,15 ± 0,575	1,726
an/ao	Region before eyes	8,3— 10,5	29,08 ± 0,738	2,214
po/ao	Region behind eyes	14,4— 21,6	49,80 ± 0,926	2,780

Tabela 2. Neki meristički karakteri hibrida *Alburnus alburnus alborella* (De Filippi, 1844) X *Leuciscus cephalus albus* (Bonaparte, 1838) iz Skadarskog jezera.

Table 2.

Some meristic characters of hybrid
Alburnus alburnus alborella (De Filippi, 1844) X
Leuciscus cephalus albus (Bonaparte, 1838) from Lake Skadar

Mark	Charakters	Amplitude (min.—max.)	M
SP.BR.	Number of the gill rakers	10—18	12,60
L.L.	Number of scales in the laterl line	46—53	49,37
	Number of scales above lateral line	8	8,00
	Number of scales under lateral line	3	3,00
D	Number of the dorsal rays	III 8	III 8,00
A	Number of the anal rays	III 9—III 14	III 10,50
T.T.	Thront teeth	2.5— 5.2	2.5—5.2

The dorstal fin of hybrids is mainly placed towards to back than that of *Leuciscus cephalus albus*. The vertical of anterior margin of base of dorsal fin is presented remarkable behind the posterior margin of ventral fin. Based on this charakters the hybrids individuals are intermediate between parents species. Behind of base ventral fin apper one subrounded keel, very characteristic

for species of *Alburnus alburnus alborella*. The keel behind base of ventral fin is sometimes less developed.

Some meristic characters of hybrids and their parents are given on table 3.

Table 3.

Mean value of some meristic characters of *Alburnus alburnus alborella* (De Filippi, 1844), *Leuciscus cephalus albus* (Bonaparte, 1838) and their hybrids

Tabela 3.

Srednja vrijednost nekih merističkih karaktera *Alburnus alburnus alborella* (De Filippi, 1844), *Leuciscus cephalus albus* (Bonaparte, 1838) i njihovih hibrida

Charakters Karaktereri	<i>Alburnus alburnus alborella</i>	Hybrid	<i>Leuciscus ce- phalus albus</i>
sp.br.	20,30	12,60	10,50
l.l.	50,28	49,37	44,65
Above l.l.	8,75	8,00	7,00
Under l.l.	3,00	3,00	3,00
D	III 7,98	III 8,00	III 8,00
A	III 13,58	III 10,50	III 8,000
Thront teeth	2.5.—5.2	2.5.—5.2	2.5.—5.2

The hybridies differ from parents by the number of branchiospines on the first gill raker (table 1).

Middle value of branchiospina of hybrids is 12.60, parents: *Alburnus alborella* 20,30 and *Leuciscus cephalus albus* 10,50. That character of hybrids is also between both parents species.

The lateral line of hybridies appear always 8 horizontal scales rows by *Alburnus alburnus alborella* average 8.75 and *Leuciscus cephalus albus* 7,0. Under lateral line of hybrids appear stable number of scales like those their parents species.

The lateral line of hybrids possesses usually 49,37 scales, less than those in *Alburnus alburnus alborella* 50,28 and more than those *Leuciscus cephalus albus* 44,65.

The hybrids and parents possess the stable number of the unbranched rays in the dorsal fin while the number of branched rays of *Alburnus alburnus alborella* is something variable. Very interesting is number of the branched rays in the anal fin of hybrids 10.50, remarkable less than in *Alburnus alburnus alborella* 13,58 but much parents species. The hybridies possess bi-thront teeth like their parents.

There are differences in position of mouth. In some hybrids (caught in August 1971) the top of mouth is placed in the upper third-way of the eye. In hybrids caught in July 1972. the top of mouth is over of horizontal margin of eye-socket.

Recently a number of interspecific hybrid forms from the family of Cyprinidae from Adriatic drainage basin in Yugoslavia have been described. In most cases the question is about the hybrid forms of endemic species so far unknown in the scientific literature. The following hybrids have been found in the rivers which disappear into the karst ground (*Phoxinus phoxinus* X *Barbus meridionalis petenyi*, Vuković, 1963), *Scardinius erythrophthalmus* X *Leuciscus turskyi*, Vuković, T. 1964, *Chodostroma phoxinus* X *Paraphoxinus alepidotus*, Vuković et al 1970), *Scardinius erythrophthalmus* X *Paraphoxinus alepidotus*, Vuković, T. et al 1971) and in the Neretva river (*Scardinius erythrophthalmus* X *Alburnus alburnus* Vuković, T. 1977, *Rutilus rubilio* X *Alburnus alburnus alborella*, Vuković, T. 1968).

The description of the hybrid between *Alburnus alburnus alborella* and *Leuciscus cephalus albus*, mentioned in this study, supplements the list of the known interspecific hybrids forms from the family of Cyprinidae from the waters of the Adriatic drainage basin in Yugoslavia. The large material on natural and artificial hybridization of fish has been gathered all over the world (Hubbs, C. L., Schultz, R. J., Nikoljukin, N. I., Schwartz, J. F. and others).

These data have a fundamental scientific importance. For the hybrid identification, as it is the case in this study, we usually use the morphological methods. However it is felt for the need a more complex understanding of the hybrids and hybridization in general, what but demands the use of other methods (cytological, physiological, biochemical etc). Therefore the information given in this paper, about the hybrids between *Alburnus alburnus alborella* and *Leuciscus cephalus albus* should be considered as a preliminary description, the first identification which has to be confirmed and supplemented with new and more complex data.

CONCLUSIONS

The hybrid forms between *Alburnus alburnus alborella* and *Leuciscus cephalus albus* exist in Lake Skadar.

These hybrids are characterized by the specific morphological features which differ them from the both parent species. It are: number of barhiospina the first gill raker, number of scales in lateral line, number of scales above lateral line, number of the branched rays in the anal fin.

LITERATURA

- Hubbs, C. L. (1955): Hybridization between fish species in nature; systematic zoology 4:1—20. In D. A. Levin editor 1979, Hybridization; an evolutionary perspective, p. 139—158 Dowder, Hutchison, and Ross, Stroudsburg, Pa., Alsoref. 696 in Publ. Gulf. Coast Res. Lab. Mus. 3, 328 p. 1972.

- Ivanović, B. (1967): Hibridizacija *Pachychilon pictum* X *Rutilus rubilio*. Polj. i šum., 13(1—2):13—18.
- Schultz, R. J. (1973): Origin and synthesis of a unisexual In: J. H. Schroder (editor), Genetics and mutagenesis of fish, p. 207—211. Springer—Verlag, Berl.
- Schwartz, F. J. (1981): World literature to Fish Hybrids with an Analysis by Family, Species, and Hybrid: Supplement 1. Gulf. Coast Research Laboratory Museum.
- Vuković, T. Ivanović, B. (1971): Slatkovodne ribe Jugoslavije. Sarajevo.
- Vuković, T. (1963): Nalaz križanca gagice i sapače (*Phoxinus phoxinus* Linnaeus X *Barbus meridionalis petenyi* Heckel (u okolini Banja Luke. Ribarstvo Jugoslavije, 18 (5).
- Vuković, T. (1964): Prilog poznavanju prirodne hibridizacije ciprinida u vodama Livanjskog polja. God. Biol. Inst. Univer. u Sarajevu, god. XVII; 199—206.
- Vuković, T. (1968): Mlaz hibrida *Rutilus rubilio* Bonaparte X *Alburnus alburnus alborella* (Filippi) u slivu rijeke Neretve. Glas. Zem. muzeja B i H, sv. VII — Prirodne nauke.
- Vuković, T., Seratlić—Savić, D., Karanac, V. (1971): Opis novog hibrida *Scardinius erythrophthalmus* X *Paraphoxinus alepidotus*. Ichthyologia, 3. (1.): 79—88.
- Vuković, T. Seratlić—Savić, D., Karanac, V. (1970): Neke morfološke karakteristike hibrida *Chodrostoma phoxinus* (Heckel) X *Paraphoxinus alepidotus* (Heckel). Ichthyologia, 2. (1.): 155—169.
- Vuković, T. (1977): Ribe Bosne i Hercegovine. Sarajevo.
- Nikoljukin, N. I. (1971): Fundamentals of hybridization in fish culture. In Seminar (study tour in the USSR. on genetic selection and hybridization of cultivated fishes, 19 April — 29 May 1968, p. 321—327. Fao UNDP/TA Rep. 2926.

MORFOLOŠKA ANALIZA NEKIH HIBRIDNIH OBLIKA RIBA IZ SKADARSKOG JEZERA

Tihomir VUKOVIĆ, Božina IVANOVIĆ, Borivoj KNEŽEVIĆ

Re z i m e

U Skadarskom jezeru egzistiraju hibridi između *Alburnus alburnus alborella* i *Leuciscus cephalus albus*. Ti hibridi se odlikuju specifičnim morfološkim osobinama po kojima se razlikuju od obje roditeljske vrste, kao što su: broj branhiospina na prvom škržnom luku, broj krljušti u bočnoj liniji, broj krljušti iznad bočne linije, broj granatih zrakova u analnom peraju.